

Joint event on

2ND ANNUAL CONGRESS ON DIABETES AND ITS COMPLICATIONS &
8TH ANNUAL CONGRESS ON PROBIOTICS, FUNCTIONAL FOODS & NUTRACEUTICALS

March 25-26, 2019 Hong Kong

Serum lipid profiles in patients with chronic kidney disease in a Saudi population

Khalid Salim Aljabri¹, Samia A Bokhari¹, Muneera A Alshareef¹, Patan M Khan¹, Hesham M Ebu Elsaoud¹, Mohammad M Jalal¹, Rania F Safwat¹, Rehab El Boraie¹ and Bandari K Aljabri²

¹King Fahad Armed Forces Hospital, Saudi Arabia

²Um Al Qura University, Saudi Arabia

Introduction & Aim: Chronic Kidney Disease (CKD) is an increasing public health issue. Serum lipids might be independent risk factors for CKD. The aim is to examine the association of serum lipid profiles, lipid ratios with CKD in a Saudi population.

Method: Three thousand six hundred and two (3602) participants who are between the age 18 to 98 years were analyzed. All patients were from the population of the diabetic center and primary health center at King Fahad Armed Forces Hospital, Jeddah, Saudi Arabia.

Results: A total of 3602 subjects (the mean age was 49.1 ± 14.1 years, minimum 18 years and maximum 98 years) were included in the analysis. 1358 (37.7%) subjects were men and 700 (19.4%) subjects had CKD. 391 (28.8%) cases were male and 309 (13.8%) cases were female with male to female ratio 2.1:1, $P < 0.0001$. Patients with CKD were significantly older than patients without CKD, (57.1 ± 12.9 vs. 47.2 ± 14.1 respectively, $p < 0.0001$). Mean BMI was significantly higher in patients compared to those patients without CKD (31.6 ± 6.4 vs. 30.8 ± 6.9 respectively, $p = 0.006$). Moreover, patients with CKD have significantly higher prevalence of type-2 diabetes and hypertension than patients without CKD (85.5% and 64.9% respectively, $p < 0.0001$). Patients with CKD have significantly higher mean and percentage of TC, LDL, TG and low HDL than patients without CKD, $p < 0.0001$. Patients with CKD had a higher TC/HDL ratio, LDL/HDL ratio and TG/HDL ratio, than patients without CKD and all of these differences were significant ($p < 0.0001$). Male had significant higher serum TC, lower LDL and lower HDL than female ($p < 0.0001$). Male had also a higher significant TG/HDL ratio, a higher TC/HDL ratio and nonsignificant a higher LDL/HDL ratio than female. Percentage of TC, LDL, TG and low HDL CKD is non-statistically consistently increasing with age.

Conclusion: Dyslipidemia either TC, LDL, TG and low HDL is high in patients with CKD. Moreover, TC/HDL, LDL/HDL and TG/HDL ratios were significantly high in patients with CKD. Thus, confirming presence of atherogenic lipid profile needing early intervention to prevent cardiovascular complications.

khalidsaljabri@yahoo.com